arXiv:1011.2049v1 [math.SP] 9 Nov 2010

Connectivity and Minimal Distance Spectral Radius of Graphs

Xiaoling Zhang[ab] [*] [†], Chris Godsil[a*]

[a]Combinatorics and Optimisation University of Waterloo,

Waterloo, Ontario, Canada, N2L 3G1

[b]School of Mathematics and Statistics, Lanzhou University,

Lanzhou, Gansu 730000, P. R. China

Abstract

In this paper, we study how the distance spectral radius behaves when the graph is perturbed by grafting edges. As applications, we also determine the graph with k cut vertices (respectively, k cut edges) with the minimal distance spectral radius.

Key words: Distance spectral radius; Pendant path

AMS subject classification: 05C50; 15A18

1 Introduction

The distance matrix of a graph, while not as common as the more familiar adjacency matrix, has nevertheless come up in several different areas, in-

---

[*] E-mail addresses: zhangxling04@lzu.cn, cgodsil@uwaterloo.ca.

[†] The author is currently a visiting Ph.D. student at the Department of Combinatorics and Optimisation in University of Waterloo from September 2008 to September 2009. She is partially supported by NSFC grant no. 10831001.

cluding communication network design [6], graph embedding theory [4, 7, 8], molecular stability [10, 11], and network flow algorithms [3, 5]. So it is interesting to study the spectra of these matrices. In this paper, we study the largest eigenvalue of the distance matrix of a graph.

Throughout this paper, we will assume that G is a simple, connected graph of order n, that is, with n vertices. Let G be a connected graph with vertex set $\{1, \ldots, n\}$. The distance between vertices i and j of G, denoted by $\mathrm{dist}(i, j)$, is defined to be the length (i.e., the number of edges) of the shortest path from i to j [2]. The distance matrix of G, denoted by $D(G)$ is the $n \times n$ matrix with its (i, j)-entry equal to $\mathrm{dist}(i, j)$, $i, j = 1, 2, \ldots, n$. Note that $\mathrm{dist}(i, i) = 0$, $i = 1, 2, \ldots, n$. The distance eigenvalue of largest magnitude is called the distance spectral radius, and is denoted by Λ_1. Balaban et al. [1] proposed the use of Λ_1 as a structure-descriptor, and it was successfully used to make inferences about the extent of branching and boiling points of alkanes[1, 9].

In this paper, we determine the graph with k cut vertices (respectively, k cut edges) which has the minimal distance spectral radius.

Main results

Let G be a connected graph. Let $\deg(v)$ (or $\deg_G(u)$) denote the degree of the vertex v in G. We define a pendant path of G to be a walk $v_0 v_1 \cdots v_s$ $(s \geqslant 1)$ such that the vertices $v_0, v_1, \ldots, v_s$ are distinct, $\deg(v_0) > 2$, $\deg(v_s) = 1$, and $\deg(v_i) = 2$, whenever $0 < i < s$. And v_0, s are called the root and the length of the pendant path, respectively.

We give a generalization of Theorem 3.5 in [12].

Theorem 1.1. *Let u and v be two adjacent vertices of a connected graph G and for positive integers k and l, let $G_{k,l}$ denote the graph obtained from G by adding paths of length k at u and length l at v. If $k > l \geqslant 1$, then $\Lambda_1(G_{k,l}) < \Lambda_1(G_{k+1,l-1})$; if $k = l \geqslant 1$, then $\Lambda_1(G_{k,l}) < \Lambda_1(G_{k+1,l-1})$ or $\Lambda_1(G_{k,l}) < \Lambda_1(G_{k-1,l+1})$.*



Fig. 1. Graphs $G_{k,l}$, $G_{k+1,l-1}$, $G_{k-1,l+1}$.

Proof: Label vertices of $G_{k,l}$, $G_{k+1,l-1}$, $G_{k-1,l+1}$ as in Figure 1. We partition $V(G_{k,l})$ into $A \cup \{k+2\} \cup B \cup C \cup D \cup E$, where

$$\begin{aligned}
A &= \{1, \ldots, k+1\},\\
B &= \{k+3, \ldots, l+k+2\},\\
C &= \{j \mid \mathrm{dist}(j, k+1) + 1 = \mathrm{dist}(j, k+2)\},\\
D &= \{j \mid \mathrm{dist}(j, k+1) = \mathrm{dist}(j, k+2)\},\\
E &= \{j \mid \mathrm{dist}(j, k+1) = \mathrm{dist}(j, k+2) + 1\}.
\end{aligned}$$

When we pass from G to G', the distances within $A \cup \{k+2\} \cup B$ and within $C \cup D \cup E$ are unchanged; the distances between A and $C \cup D \cup E$, $\{k+2\}$ and E are increased by 1; the distances between B and $C \cup D \cup E$, $\{k+2\}$ and C are decreased by 1; the distances between $\{k+2\}$ and D are unchanged. If the distance matrices are partitioned according to A, $\{k+2\}$, B, C, D, E, their difference is

$$D(G_{k+1,l-1})-D(G_{k,l}) = \begin{bmatrix} 0 & 0 & 0 & e_A(e_C)^t & e_A(e_D)^t & e_A(e_E)^t \\ 0 & 0 & 0 & -(e_C)^t & 0 & (e_E)^t \\ 0 & 0 & 0 & -e_C(e_C)^t & -e_C(e_D)^t & -e_C(e_E)^t \\ (e_C)^te_A & -e_C & -(e_C)^te_C & 0 & 0 & 0 \\ (e_D)^te_A & 0 & -(e_D)^te_C & 0 & 0 & 0 \\ (e_E)^te_A & e_E & -(e_E)^te_C & 0 & 0 & 0 \end{bmatrix},$$

where

$$e_i = (\underbrace{1,\dots,1}_{|i|})^t$$

and $i = A, C, D, E$. Let $x = (x_1, \dots, x_n)^t$ be a positive eigenvector corresponding to $\Lambda_1(G_{k,l})$. Then we have

$$\begin{aligned} \frac{1}{2}\big(\Lambda_1(G_{k+1,l-1}) - \Lambda_1(G_{k,l})\big) &\geqslant \frac{1}{2}x^t\big(D(G_{k+1,l-1}) - D(G_{k,l})\big)x \qquad (1.1) \\ &= \left(\sum_{j\in A} x_j - \sum_{j\in B} x_j\right) \sum_{j\in C\cup D\cup E} x_j \\ &\quad -x_{k+2}\sum_{j\in C} x_j + x_{k+2}\sum_{j\in E} x_j \\ &\geqslant \left(\sum_{j\in A} x_j - \sum_{j\in B} x_j - x_{k+2}\right) \sum_{j\in C\cup D\cup E} x_j. \end{aligned}$$

Similarly, we have

$$\begin{aligned} \frac{1}{2}\big(\Lambda_1(G_{k-1,l+1}) - \Lambda_1(G_{k,l})\big) &\geqslant \frac{1}{2}x^t\big(D(G_{k-1,l+1})) - D(G_{k,l})\big)x \\ &= \left(\sum_{j\in B} x_j + x_{k+2} - \sum_{j\in A} x_j + x_{k+1}\right) \sum_{j\in C\cup D\cup E} x_j \\ &\quad +x_{k+1}\sum_{j\in C} x_j - x_{k-2}\sum_{j\in E} x_j \\ &\geqslant \left(\sum_{j\in B} x_j + x_{k+2} - \sum_{j\in A} x_j\right) \sum_{j\in C\cup D\cup E} x_j. \end{aligned}$$

Since $D(G_{k,l})x = \Lambda_1(G_{k,l})x$, i.e., $\Lambda_1(G_{k,l})x_i = \sum\limits_{j=1}^{n} \mathrm{dist}(i,j)x_j$, for $1 \leqslant i \leqslant n$, we get:

if $1 \leqslant i \leqslant l+1$, then

$$\begin{aligned}\Lambda_1(G_{k,l})(x_i - x_{l+k+3-i}) &= \sum_{m=1}^{s}(l+k-2m+3)(x_{l+k+3-m} - x_m) + (k-l)\sum_{j\in D} x_j \\ &\quad +(k-l-1)\sum_{j\in C} x_j + (k-l+1)\sum_{j\in E} x_j \\ &\quad -2\sum_{m=1}^{i-1}(x_{l+k+3-m} - x_m);\end{aligned} \tag{1.2}$$

if $l+2 \leqslant i \leqslant s$, then

$$\begin{aligned}\Lambda_1(G_{k,l})(x_i - x_{l+k+3-i}) &= \sum_{m=1}^{s}(l+k-2m+3)(x_{l+k+3-m} - x_m) \\ &\quad +(k+l+3-2i)\sum_{j\in C\cup D\cup E} x_j - 2\sum_{m=1}^{i-1}(x_{l+k+3-m} - x_m);\end{aligned} \tag{1.3}$$

where $s = \frac{l+k+2}{2}$, if $l+k$ is even; and $s = \frac{l+k+1}{2}$, if $l+k$ is odd.

From Eqs. (1.3) and (1.4), we get:

if $2 \leqslant i \leqslant l+1$, then

$$\begin{aligned}&\Lambda_1(G_{k,l})(x_{i-1} - x_{l+k-i+4}) - \Lambda_1(G_{k,l})(x_i - x_{l+k+3-i}) \\ &= 2\sum_{r=1}^{i-1}(x_{l+k+3-r} - x_r);\end{aligned} \tag{1.4}$$

if $i = l+2$, then

$$\begin{aligned}&\Lambda_1(G_{k,l})(x_{i-1} - x_{l+k-i+4}) - \Lambda_1(G_{k,l})(x_i - x_{l+k+3-i}) \\ &= 2\sum_{r=1}^{i-1}(x_{l+k+3-r} - x_r) + \sum_{j\in D} x_j + 2\sum_{j\in E} x_j;\end{aligned} \tag{1.5}$$

if $l+3 \leqslant i \leqslant s$, then

$$\begin{aligned}&\Lambda_1(G_{k,l})(x_{i-1} - x_{l+k-i+4}) - \Lambda_1(G_{k,l})(x_i - x_{l+k+3-i}) \\ &= 2\sum_{r=1}^{i-1}(x_{l+k+3-r} - x_r) + 2\sum_{j\in C\cup D\cup E} x_j.\end{aligned} \tag{1.6}$$

Equation (1.5) implies that, for $1 \leqslant i \leqslant l+1$, the differences $x_{l+k+3-i} - x_i$ are either all non-positive or non-negative.

Case 1. $k = l$.

If $x_{l+k+3-i} - x_i \leqslant 0$, for all $1 \leqslant i \leqslant l+1$, then

$$\sum_{j\in A} x_j - \sum_{j\in B} x_j - x_{k+2} \geqslant 0.$$

So we get $\Lambda_1(G_{k,l}) \leqslant \Lambda_1(G_{k+1,l-1})$. Here, if $\Lambda_1(G_{k,l}) = \Lambda_1(G_{k+1,l-1})$, then the last equality in (1.2) holds, which implies that

$$2\sum_{j\in E} x_j + \sum_{j\in D} x_j = 0,$$

i.e., $D = E = \emptyset$. Since $i-1 \leqslant l = s-1$, for $1 \leqslant m \leqslant i-1$, $l+k-2m+3 \geqslant 2$, we get

$$\sum_{m=1}^{s}(l+k-2m+3)(x_{l+k+3-m} - x_m) - 2\sum_{m=1}^{i-1}(x_{l+k+3-m} - x_m) \leqslant 0.$$

Combining this with the fact that x is a positive eigenvector corresponding to $\Lambda_1(G_{k,l})$, we get that the right side of equation (1.3) is strictly less than 0, which contradicts the fact that the left side of equation (1.3) $\Lambda_1(G_{k,l})(x_i - x_{l+k+3-i}) \geqslant 0$. So we get $\Lambda_1(G_{k,l}) < \Lambda_1(G_{k+1,l-1})$.

If $x_{l+k+3-i} - x_i \geqslant 0$, for all $1 \leqslant i \leqslant l+1$, we can get $\Lambda_1(G_{k,l}) < \Lambda_1(G_{k-1,l+1})$ similarly.

Case 2. $k > l$.

If $x_{l+k+3-i} - x_i \geqslant 0$, for all $1 \leqslant i \leqslant l+1$, then from Eqs. (1.6) and (1.7), we can get that $x_{l+k+3-i} - x_i \geqslant 0$, for all $l+2 \leqslant i \leqslant s$. Similar to the above case, we get that the right side of equation (1.5) is strict larger than 0, which contradicts the fact that the left side of equation (1.3) $\Lambda_1(G_{k,l})(x_i - x_{l+k+3-i}) \leqslant 0$. So this case is impossible.

If $x_{l+k+3-i} - x_i < 0$, then $\sum\limits_{j\in A} x_j - \sum\limits_{j\in B} x_j - x_{k+2} > 0$, which implies that

$$\Lambda_1(G_{k,l}) > \Lambda_1(G_{k+1,l-1}).$$

This completes the proof. □

From the proof of above theorem, we get the following corollary.

Corollary 1.2. *Let v_l and v_m be two adjacent vertices of connected graph G. Let P_l and P_m be two pendant paths with roots v_l and v_m, respectively. If $l > m$, then*

$$\sum_{j\in V(P_l)} x_j > \sum_{j\in V(P_m)} x_j.$$

Theorem 1.3. *Let C_1 be a component of $G-u$ and $v_1,\ldots,v_k$ $(1 \leqslant k \leqslant \deg_G(u) - \deg_{C_1}(u))$ be some vertices of $N_G(u)\setminus N_{C_1}(u)$. Suppose $N_{C_1}(u)\setminus\{v\} = N_{C_1}(v)$, where v is a vertex of C_1 adjacent to u. Let G' be the graph obtained from G by deleting the edges uv_s and adding the edges vv_s $(1 \leqslant s \leqslant k)$. If there exists a vertex $w \in V(G)\setminus(V(C_1)\cup\{u\})$ such that $\operatorname{dist}_G(w,v_s) < \operatorname{dist}_{G'}(w,v_s)$, for all $1 \leqslant s \leqslant k$, then $\Lambda_1(G) < \Lambda_1(G')$.*

Proof: We partition $V(G)$ into $A\cup B\cup\{u\}\cup\{v\}\cup C$, where

$$\begin{aligned} A &= \{v_1,\ldots,v_k\},\\ B &= V(G)\setminus(A\cup V(C_1)\cup\{u\}),\\ C &= V(C_1)\setminus\{v\}. \end{aligned}$$

From G to G', the distances between $A\cup B$ and C are unchanged; the distances between A and $\{u\}\cup\{w\}$ are increased by 1; the distances between A and $\{v\}$ are decreased by 1; the distances between A and $B\setminus\{w\}$ are not decreased. Let $x=(x_1,\ldots,x_n)^t$ be a positive eigenvector corresponding to $\Lambda_1(G)$. Similar to the proof of above theorem, we have

$$\begin{aligned} \frac{1}{2}\big(\Lambda_1(G')-\Lambda_1(G)\big) &\geqslant \frac{1}{2}x^t\big(D(G')-D(G)\big)x\\ &\geqslant (x_w+x_u-x_v)\sum_{j\in A} x_j. \end{aligned}$$

Since $D(G)x = \Lambda_1(G)x$, i.e., $\Lambda_1(G)x_i = \sum\limits_{j=1}^{n} \mathrm{dist}(i,j)x_j$, we can easily get

$$\begin{aligned}
(\Lambda_1(G)+1)(x_w+x_u-x_v) &= \sum_{j=1}^{n} (\mathrm{dist}(w,j)+\mathrm{dist}(u,j)-\mathrm{dist}(v,j))\,x_j + (x_w+x_u-x_v) \\
&= \sum_{j\in A\cup(B\setminus\{w\})} (\mathrm{dist}(w,j)+\mathrm{dist}(u,j)-\mathrm{dist}(v,j))x_j \\
&\quad +\sum_{j\in C}(\mathrm{dist}(w,j)+\mathrm{dist}(u,j)-\mathrm{dist}(v,j))x_j \\
&\quad +(\mathrm{dist}(w,w)+\mathrm{dist}(u,w)-\mathrm{dist}(v,w))x_w \\
&\quad +(\mathrm{dist}(w,u)+\mathrm{dist}(u,u)-\mathrm{dist}(v,u))x_u \\
&\quad +(\mathrm{dist}(w,v)+\mathrm{dist}(u,v)-\mathrm{dist}(v,v))x_v + (x_w+x_u-x_v).
\end{aligned}$$

As we know, $\mathrm{dist}(u,j)-\mathrm{dist}(v,j) = -1$ and $\mathrm{dist}(w,j) \geqslant 1$, for $j \in A\cup(B\setminus\{w\})$, so

$$\sum_{j\in A\cup(B\setminus\{w\})} (\mathrm{dist}(w,j)+\mathrm{dist}(u,j)-\mathrm{dist}(v,j))x_j \geqslant 0.$$

Similarly, we can get

$$\sum_{j\in C}(\mathrm{dist}(w,j)+\mathrm{dist}(u,j)-\mathrm{dist}(v,j))x_j > 0.$$

Since

$$\begin{aligned}
\mathrm{dist}(w,w)+\mathrm{dist}(u,w)-\mathrm{dist}(v,w) &= -1, \\
\mathrm{dist}(w,u)+\mathrm{dist}(u,u)-\mathrm{dist}(v,u) &\geqslant 0, \\
\mathrm{dist}(w,v)+\mathrm{dist}(u,v)-\mathrm{dist}(v,v) &\geqslant 3,
\end{aligned}$$

combining these with all the above equations and inequations, we get

$$\big(\Lambda_1(G)+1)(x_w+x_u-x_v\big) > 0,$$

which implies $\Lambda_1(G') > \Lambda_1(G)$.

This completes the proof. $\square$

2 Applications

The graph $G_{n,k}$ is a graph obtained by adding paths $P_{l_1+1}, \ldots, P_{l_{n-k}+1}$ of almost equal lengths (by the length of a path, we mean the number of its vertices) to the vertices of the complete graph K_{n-k}; that is, the lengths $l_1, \ldots, l_{n-k}$ of $P_{l_1+1}, \ldots, P_{l_{n-k}+1}$ which satisfy $|l_i - l_j| \leqslant 1$; $1 \leqslant i, j \leqslant n-k$.

K_n^k is a graph obtained by joining k independant vertices to one vertex of K_{n-k}.

Theorem 2.1. *Of all the connected graphs with n vertices and k cut vertices, the minimal distance spectral radius is obtained uniquely at $G_{n,k}$.*

Proof: We are supposed to prove that if G is a connected graph with n vertices and k cut vertices, then $\Lambda_1(G) \geqslant \Lambda_1(G_{n,k})$ with equality only when $G \cong G_{n,k}$. Let V_1 be the set of the cut vertices of G. Note that if we add some edges to G such that each block of $G - V_1$ is a clique, denoting the new graph by G', then $\mathrm{dist}_G(i,j) \geqslant \mathrm{dist}_{G'}(i,j)$. Consequently, $D(G) \geqslant D(G')$, which implies $\Lambda_1(D(G)) \geqslant \Lambda_1(D(G'))$ with equality only when $D(G) = D(G')$. So, in the following, we always assume that each cut vertex of G connects exactly two blocks and that all these blocks are cliques. Order the cardinalities of these blocks $a_1 \geqslant a_2 \geqslant \cdots \geqslant a_{k+1} \geqslant 2$ and denote the blocks by $K_{a_1}, \ldots, K_{a_{k+1}}$. If $k = 0$, then $G \cong K_n$ and the theorem holds. If $k = n-2$, then G is the path $G_{n,n-2}$. If $k = n-3$, then $a_1 = 3$, $a_2 = \cdots = a_{k+1} = 2$. The result follows from a repeated use of Theorem 1.1. Thus we may assume that $1 \leqslant k \leqslant n-4$. Moreover, we observe that $a_1 = n+k-(a_2+\cdots+a_{k+1}) \leqslant n-k$.

Choose G such that the distance spectral radius is as small as possible.

Claim. $a_1 = n - k$.

Otherwise, $a_1 \leqslant n-k-1$, which implies $a_2 \geqslant \cdots \geqslant a_i \geqslant 3$ for some i, $2 \leqslant i \leqslant k+1$.

Suppose $K_{a_{i_1}}, \ldots, K_{a_{i_t}}$ are the blocks, each of which contains at least two roots of pendant paths of G. Let $\mathcal{P}$ be the set of pendant paths whose roots are contained in $K_{a_{i_1}}, \ldots, K_{a_{i_t}}$ and P_m be one of the shortest pendant paths among $\mathcal{P}$. Suppose the root of P_m is contained in $K_{a_{i_s}}$ for some $1 \leqslant s \leqslant t$

and P_l is another pendant path whose root is also contained in $K_{a_{i_s}}$. Then we have $0 \leqslant l - m \leqslant 1$. Otherwise, by Theorem 1.1, we can find a graph G' such that $\Lambda_1(G) > \Lambda_1(G')$, which is a contradiction. We label the vertices of G such that

$$\begin{aligned} P_m &= v_1 \cdots v_m, \\ P_l &= v_{m+1} \cdots v_{m+l}, \\ V(K_{a_{i_s}}) \setminus \{v_m, v_{m+1}\} &= \{u_1, \ldots, u_r\}, \end{aligned}$$

where v_m and v_{m+1} are the roots of P_m and P_l, respectively. Suppose u_1 is a cut vertex of G such that $G[V(C_1) \cup \{u_1\}]$ contains at least one block K_{a_h}, for some $1 \leqslant h \leqslant i$, where C_1 is the component of $G - u_1$ which does not contain v_m.

Let $N_{C_1}(u_1) = \{w_1. \ldots, w_q\}$ and

$$\begin{aligned} G' = G &- v_m u_2 - \cdots - v_m u_r - v_m v_{m+1} \\ &+ w_1 u_2 + \cdots + w_1 u_r + w_1 v_{m+1} + \\ &\cdots\cdots\cdots \\ &+ w_{q-1} u_2 + \cdots + w_{q-1} u_r + w_{q-1} v_{m+1} \\ &+ w_q u_2 + \cdots + w_q u_r + w_q v_{m+1}. \end{aligned}$$

Then G' is a connected graph with n vertices and k cut vertices.

In the following, we consider the graph G'. Since $V(G') = V(G)$, we partition $V(G')$ into $A \cup \{u_1\} \cup B \cup C$, where

$$\begin{aligned} A &= V(P_m), \\ B &= V(C_1), \\ C &= V(G) \setminus (A \cup \{u_1\} \cup B). \end{aligned}$$

From G' to G, the distances between A and $B \cup \{u_1\}$, $\{u_1\}$ and B are unchanged; the distances between B and C are increased by 1; the distances between A and C are decreased by 1. Let $x = (x_1, \ldots, x_n)^t$ be a positive eigenvector corresponding to $\Lambda_1(G')$. Then we have

$$\begin{aligned}\frac{1}{2}\big(\Lambda_1(G)-\Lambda_1(G')\big) &\geqslant \frac{1}{2}x^t\big(D(G)-D(G')\big)x \\ &\geqslant \left(\sum_{j\in B}x_j-\sum_{j\in A}x_j\right)\sum_{j\in C}x_j.\end{aligned} \tag{2.1}$$

Case 1. $\sum\limits_{j\in B}x_j-\sum\limits_{j\in A}x_j>0$.

From inequality (2.2), we get that $\Lambda_1(G)>\Lambda_1(G')$, which is a contradiction.

Case 2. $\sum\limits_{j\in B}x_j-\sum\limits_{j\in A}x_j\leqslant 0$

Since $D(G')x=\Lambda_1(G')x$, i.e., $\Lambda_1(G')x_i=\sum\limits_{j=1}^{n}\mathrm{dist}(i,j)x_j$, we can easily get that

$$\begin{aligned}\Lambda_1(G')\left(\sum_{j\in B}x_j-\sum_{j\in A}x_j\right) &= \sum_{i\in A\cup\{u_1\}}\left(\sum_{u\in B}\mathrm{dist}(u,i)-\sum_{v\in A}\mathrm{dist}(v,i)\right)x_i \\ &\quad+\sum_{i\in B}\left(\sum_{u\in B}\mathrm{dist}(u,i)-\sum_{v\in A}\mathrm{dist}(v,i)\right)x_i \\ &\quad+\sum_{i\in C}\left(\sum_{u\in B}\mathrm{dist}(u,i)-\sum_{v\in A}\mathrm{dist}(v,i)\right)x_i.\end{aligned} \tag{2.2}$$

Since $G[V(C_1)\cup\{u_1\}]=G'[V(C_1)\cup\{u_1\}]$ contains at least one block K_{a_h}, for some $1\leqslant h\leqslant i$, $G'[V(C_1)\cup\{u_1\}]$ must contain at least two pendant paths P' and P'' whose roots are contained in the same block. Denote the roots of P' and P'' by ω_1 and ω_2, respectively. Suppose $dist_{G'}(v_{m+1},\omega_1)=k$, then $k\geqslant 2$ and $dist_{G'}(v_{m+1},\omega_2)=k$. As we know, P' and P'' are two pendant paths of length at least m, so
if $i\in A\cup\{u_1\}$, then

$$\sum_{u\in B}\mathrm{dist}(u,i)-\sum_{v\in A}\mathrm{dist}(v,i)\geqslant 2\times\frac{(m+k-1)(m+k)}{2}-\frac{m(m-1)}{2}; \tag{2.3}$$

if $i \in B$, then

$$\sum_{u\in B}\mathrm{dist}(u,i)-\sum_{v\in A}\mathrm{dist}(v,i) \geqslant \frac{(k-2)(k-1)}{2}-(m-1)k; \tag{2.4}$$

if $i \in C$, then

$$\sum_{u\in B}\mathrm{dist}(u,i)-\sum_{v\in A}\mathrm{dist}(v,i) > 0. \tag{2.5}$$

is obvious.

Combining Eqs. (2.3), (2.3), (2.4) with (2.5), we get

$$\begin{aligned}
&\Lambda_1(G')\left(\sum_{j\in B}x_j-\sum_{j\in A}x_j\right)\\
&> \left[2\times\frac{(m+k-1)(m+k)}{2}-\frac{m(m-1)}{2}\right]\sum_{i\in A\cup\{u_1\}}x_i\\
&\quad+\left[\frac{(k-2)(k-1)}{2}-(m-1)k\right]\sum_{i\in B}x_i\\
&> \left[(m+k-1)(m+k)-\frac{m(m-1)}{2}+\frac{(k-2)(k-1)}{2}-(m-1)k\right]\sum_{i\in B}x_i\\
&> 0,
\end{aligned}$$

which is a contradiction. So this case does not exist.

Up to now, we have proved the claim that $a_1 = n-k$, which implies that

$$a_2=\cdots=a_{k+1}=2.$$

From a repeated use of Theorem 1.1, we get that $G \cong G_{n,k}$.

This completes the proof. $\square$

Theorem 2.2. *Of all the connected graphs with n ($n \geqslant 4$) vertices and k cut edges, the minimal distance spectral radius is obtained uniquely at K_n^k.*

Proof: We are supposed to prove that if G is a connected graph with n vertices and k cut edges, then $\Lambda_1(G) \geqslant \Lambda_1(K_n^k)$ with equality only when

$G \cong K_n^k$. Let $E_1 = \{e_1, e_2, \ldots, e_k\}$ be the set of the cut edges of G. For a similar reason to Theorem 2.1, we assume that each component of $G - E_1$ is a clique.

If $k = 0$, then $G \cong K_n$ and the theorem holds. So we assume that $k \geqslant 1$. Denote the components of $G - E_1$ by $K_{a_0}, \ldots, K_{a_k}$, where $a_0 + \cdots + a_k = n$.

Let $V_{a_i} = \{v \in K_{a_i} : v \text{ is an end vertex of the cut edges of } G\}$, and choose G such that the distance spectral radius is as small as possible.

Claim 1. $|V_{a_i}| = 1$, $0 \leqslant i \leqslant k$.

Otherwise, $|V_{a_i}| > 1$ for some $0 \leqslant i \leqslant k$. Suppose u, $v \in V_{a_i}$ and $vv_j, uv_h \in E_1$. Let

$$G' = G - vv_j + uv_j.$$

Then, G' is still a connected graph with n vertices and k cut edges, and $\mathrm{dist}_{G'}(v_h, v_j) < \mathrm{dist}_G(v_h, v_j)$. In G', $N_{K_{a_i}}(u) \setminus \{v\} = N_{K_{a_i}}(v) \setminus \{u\}$, so by Theorem 1.3, we get $\Lambda_1(G') < \Lambda_1(G)$, which is a contradiction.

So, in the following, we can assume that $V_{a_i} = \{v_i\}$, $0 \leqslant i \leqslant k$.

Claim 2. If v_s is adjacent to v_t, where $0 \leqslant t, s \leqslant k$, then $\deg(v_s) = 1$ or $\deg(v_t) = 1$.

Otherwise, $\deg(v_s) \geqslant 2$ and $\deg(v_t) \geqslant 2$. Denote by C_1, C_2 the components of $G - v_s v_t$ which contain v_s and v_t, respectively. Suppose $N(v_s) \setminus \{v_t\} = \{v_{k+1}, \ldots, v_{k+r}\}$ and $u \in N(v_t) \setminus \{v_s\}$. Let

$$G' = G - v_s v_{k+1} - \ldots - v_s v_{k+r} + v_t v_{k+1} + \ldots + v_t v_{k+r}.$$

Then, G' is still a connected graph with n vertices and k cut edges, and $\mathrm{dist}_{G'}(u, v_{k+i}) < \mathrm{dist}_G(u, v_{k+i})$ $(1 \leqslant i \leqslant r)$. In G', $v_s v_t$ is a pendant edge, so by Theorem 1.3, we get $\Lambda_1(G') < \Lambda_1(G)$, which is a contradiction.

Since G is connected, combining Claim 1 with Claim 2, we get that $G \cong K_n^k$.

This completes the proof. $\square$

References

[1] A. T. Balaban, D. Ciubotariu, M. Medeleanu, Topological indices and real number vertex invariants based on graph eigenvalues or eigenvectors. J. Chem. Inf. Comput. Sci. 31 (1991) 517-523.

[2] F. Buckley, F. Harary, Distance in Graphs, Addison-Wesley, Redwood, 1990.

[3] E. W. Dijkstra, A note on two problems in connection with graphs. Numer. Math. 1 (1959) 269-271.

[4] M. Edelberg, M. R. Garey and R. L. Graham, On the distance matrix of a tree. Discrete Math. 14 (1976) 23-29.

[5] M. L. Fredman, New bounds on the complexity of the shortest path problem. SIAM J. Comput. 5 (1976) 83-89.

[6] R. L. Graham and H. O. Pollak, On the addressing problem for loop switching. Bell Sys. Tech. J. 50 (1971) 2495-2519.

[7] R. L. Graham and H. O. Pollak, On embedding graphs in squashed cubes. Graph Theory ond Applications. Springer, Berlin (1973) 99-110.

[8] R. L. Graham and L. Lovasz, Distance matrix polynomials of trees. Adv. in Math. 29 (1978) 60–88.

[9] I. Gutman, M. Medeleanu, On structure-dependence of the largest eigenvalue of the distance matrix of an alkane, Indian J. Chem. A 37 (1998) 569-573.

[10] H. Hosoya, M. Murakami and M. Gotoh, Distance polynomial and characterization of a graph. Natur. Sci. Rep. Ochanomizu Univ. 24 (1973) 27-34.

[11] D. H. Rouvray, The search for useful topological indices in chemistry. Amer. Scientist 61 (1973) 729-735.

[12] S. N. Ruzieh, D. L. Powers, The distance spectrum of the path P_n and the first distance eigenvector of connected graphs, Linear and Multilinear Algebra, 28:1 (1990) 75-81.